ELSEVIER

Fluid Phase Equilibria 110 (1995) ix–x

FLUID PHASE EQUILIBRIA

Contents

(Abstracted / Indexed in: Curr. Contents / Eng. Technol. Appl. Sci. Curr. Contents / Phys. Chem. Earth Sci., Sci. Cit. Index, Phys. Abstr., ASCA, Geo Abstracts, GEOBASE)

Theoretical papers

Thermodynamic study of the micellar solution – solid surfactant equilibrium
N.A. Smirnova (St. Petersburg, Russia) 1–15

Phase equilibria modeling by the quasilattice equation of state for binary and ternary systems composed of carbon dioxide, water and some organic components
I.V. Prikhodko, A.I. Victorov, N.A. Smirnova (St. Petersburg, Russia) and Th.W. de Loos (Delft, The Netherlands) 17–30

Analysis of the intramolecular proximity effect on dichloroalkane + alkane mixtures using Nitta–Chao model
J. García, E.R. López (Santiago de Compostela, Spain), J.L. Legido (Vigo, Spain) and J. Fernández (Santiago de Compostela, Spain) 31–51

Estimation of the Nitta–Chao parameter for the ether-group. Ether + *n*-alkane mixtures
J. Vijande, J.L. Legido, T.P. Iglesias (Vigo, Spain), E. López and J. Fernández (Santiago de Compostela, Spain) 53–71

Thermodynamics of systems of (oxy)alcohols and (poly)ethers
D. Missopolinou and C. Panayiotou (Thessaloniki, Greece) 73–88

Thermodynamic consistency of data banks
D. Miklós, S. Kemény, G. Almásy and K. Kollár-Hunek (Budapest, Hungary) 89–113

Experimental papers

Measurement and correlation of partition coefficients of hydrolytic enzymes for dextran + poly (ethylene glycol) + water aqueous two-phase systems at 20°C
T. Furuya, Y. Iwai, Y. Tanaka, H. Uchida, S. Yamada and Y. Arai (Fukuoka, Japan) 115–128

Langmuir films of mixtures of C_{70} and arachidic acid
R. Castillo, S. Ramos (México, México), J. Ruíz-García (San Luis Postosí, México) and C. Varea (México, México) . . . 129–135

Water transfer in mixed water-in-oil emulsions studied by differential scanning calorimetry
D. Clausse, I. Pezron and A. Gauthier (Compiègne cedex, France) 137–150

Fluid phase equilibria of binary and ternary mixtures of supercritical carbon dioxide with tetradecanoic acid and docosane up to 43 MPa and 393 K: cosolvency effect and miscibility windows
H. Pöhler and G.M. Schneider (Bochum, Germany) 151–156

Fluid multiphase equilibria and critical phenomena in binary and ternary mixtures of carbon dioxide, certain n-alkanols and tetradecane
C.J. Peters, L.J. Florusse, S. Hähre and J. de Swaan Arons (Delft, The Netherlands) 157–173

Vapour–liquid equilibrium at high pressure in the system containing carbon dioxide and propyl acetate
Z. Wagner (Praha, Czech Republic) 175–182

Phase equilibria in binary systems containing N-monosubstituted amides and hydrocarbons
J. Schmelzer and J. Pusch (Leipzig, Germany) 183–196

Measurement and correlation of liquid–liquid equilibria of methanol + alkane + ether ternary systems
H. Higashiuchi, Y. Sakuragi (Kurume, Japan) and Y. Arai (Fukuoka, Japan) 197–204

Isobaric vapor-liquid equilibria of acetone–propyl ether and isopropyl ether–propyl ether systems. Corroboration of no reverse volatility
J.M. Resa, M.A. Betolaza, C. González and A. Ruiz (Vitoria, Spain) 205–217

Vapor–liquid equilibria of binary mixtures containing methyl tert-butyl ether (MTBE) and/or substitution hydrocarbons at 298.15 K and 313.15 K
L.M. Lozano, E.A. Montero, M.C. Martín and M.A. Villamañán (Valladolid, Spain) 219–230

Thermodynamics of the ternary mixture propan-1-ol + tetrahydrofuran + n-heptane at 298.15 K. Experimental results and ERAS model calculations of G^E, H^E and V^E
M. Keller, S. Schnabel and A. Heintz (Heidelberg, Germany) 231–265
Thermodynamic properties (H_m^E, $C_{P,m}^E$, V_m^E, κ_T^E) of binary mixtures{x1,3-Dioxane + $(1-x)$Cyclohexane} at 298.15 K
T. Takigawa (Osaka, Japan), H. Ogawa (Saitama, Japan), M. Nakamura, K. Tamura and S. Murakami (Osaka, Japan) . . . 267–281
Thermodynamics of mixtures formed by polycyclic aromatic hydrocarbons with long chain alkanes
A. Aoulmi, M. Bouroukba, R. Solimando (Nancy Cédex, France) and M. Rogalski (Metz, France) 283–297
Thermodynamics of (1,4-difluorobenzene + an n-alkane) and of (hexafluorobenzene + an n-alkane)
E. Wilhelm (Wien, Austria), A. Inglese (Bari, Italy), A. Lainez (Madrid, Spain), A.H. Roux and J.-P.E. Grolier (Aubière, France) 299–313
Determination of the average molecular weight of petroleum cuts by vapor pressure depression
A. Blondel-Telouk, H. Loiseleur (Villeurbanne, France), A. Barreau, E. Behar (Rueil-Malmaison, France) and J. Jose (Villeurbanne, France) 315–339
Thermophysical properties of 1-hexanol over the temperature range from 303 K to 503 K and at pressures from the saturation line to 400 MPa
S.L. Randzio (Warszawa, Poland), J.-P.E. Grolier and J.R. Quint (Aubière, France) 341–359
Excess molar volumes of 1-alcohol + aliphatic monoethers at 298.15 K
A. Serna, I. García de la Fuente, J.A. González, J.C. Cobos and C. Casanova (Valladolid, Spain) 361–367
Densities and viscosities of the binary mixtures of interest for absorption refrigeration systems and heat pumps
X. Esteve, F. Olive, K.R. Patil, S.K. Chaudhari and A. Coronas (Tarragona, Spain) 369–382

Subject Index 383–384

Author Index 385